

Diagnosis	No.	Age (<1 year)	Age (1-5 years)	Age (>5 year)	P value
Complicated Pneumonia	11	9 (81.8%)	2 (18.2%)	0	<0.0001
Complicated CHD	8	3 (37.5%)	4 (50.0%)	1 (12.5%)	
DKA	7	0	3 (42.9%)	4 (57.1%)	
Encephalitis	7	4 (57.1%)	3 (42.9%)	0	
GE	5	5 (100%)	0	0	
ICH	4	4 (100%)	0	0	
RF	4	0	1 (25.0%)	3 (75%)	
Scorpion bite	4	0	4 (100%)	0	
Sepsis	2	1 (50.0%)	1 (50.0%)	0	
Metabolic acidosis	2	2 (100%)	0	0	
Arrhythmia	1	1 (100%)	0	0	
cannabis toxicity	1	0	1 (100%)	0	
Organophosphorus poisoning	1	0	1 (100%)	0	
HSM	1	1 (100%)	0	0	

Table (1) Causes and risk factors by age group

Variable	Death N=25	Improved N=33	P-value
Age group			
< 1 years	16 (53.33)	14 (46.67%)	0.11
1-5 years	8 (40.00%)	12 (60.00%)	
>5 years	1 (12.50%)	7 (87.50%)	
Sex			
Female	12 (46.15%)	14 (53.85%)	0.67
Males	13 (40.63%)	19 (59.38%)	

Table (2) Relation between outcome and age and gender.

Variable	Death N=25	Improved N=33	P-value
Complicated Pneumonia	6 (54.55%)	5 (45.45%)	0.07
Complicated CHD	3 (37.50%)	5 (62.50%)	
DKA	0	7 (100%)	
Encephalitis	6 (85.71%)	1 (14.29%)	
GE	1 (20.00%)	4 (80.00%)	
ICH	3 (75.00%)	1 (25.00%)	
RF	1 (25.00%)	3 (75.00%)	
Scorpion bite	1 (25.00%)	3 (75.00%)	
Sepsis	2 (100%)	0	
Metabolic acidosis	1 (50.00%)	1 (50.00%)	
Arrhythmia	0	1 (100%)	
Hashish toxicity	0	1 (100%)	
Organophosphorus poisoning	0	1 (100%)	
HSM	1 (100%)	0	

Table (3) Relation between outcome and Diagnosis

	Death N=25	Improved N=33	P value
GCS Mean ± SD Median (range)	7.4±0.91 7 (6:9)	9.18±1.13 9 (7:12)	<0.0001

Table (4) Relation between outcome and GCS