

| Characteristic | N (%) |
|--|---|
| Age: Young age (18-) Middle age (40-) Old age (>65) | 40 (9.8%) 167 (40.8%) 202 (49.4%) |
| Gender: Male Female | 235 (57.5%) 174 (42.5%) |
| Marital status: Unmarried Married | 146 (35.7%) 263 (64.3%) |
| Residence Rural Urban | 243(59.5%) 166(40.5%) |
| Education level: Illiterate Primary/Secondary Higher | 144 (35.2%) 175 (42.8%) 90 (22%) |
| Occupation: Unemployed Employed | 133 (32.5%) 276 (67.5%) |
| Income Low Moderate High | 190(46.5%) 131(32%) 88(21.5%) |
| Comorbidities No Yes | 99 (24.2%) 310 (75.8%) |
| No. of antihypertensive drugs: 1 2 3-4 | 254 (62.1%) 99 (24.2%) 56 (13.7%) |
| Smoking: No Yes | 212 (51.8%) 197 (48.2%) |

Table (1): Distribution of the participants by their socio-demographic characteristics in 2018

| Characteristic | OR | 95% CI | P-value |
|---------------------------------------|-----------|---------------|----------------|
| Age: | | | |
| 18- | 7.29 | 2.36-22.53 | 0.00 |
| 40- | 2.58 | 1.48-4.5 | 0.00 |
| >65 | 1 | | 0.00 |
| Residence | | | |
| Rural | 1 | | |
| Urban | 2.93 | 1.54-4.89 | 0.03 |
| Education level: | | | 0.03 |
| Illiterate | 1 | | |
| Primary/Secondary | 2.13 | 1.19-3.8 | 0.01 |
| Higher | 2.13 | 1.02-4.45 | 0.04 |
| Income | | | 0.00 |
| Low | 1 | | |
| Moderate | 2.51 | 1.23-4.93 | 0.00 |
| High | 4.86 | 2.03-9.06 | 0.00 |
| Comorbidities: | | | |
| No | 2.25 | 1.16-4.38 | 0.02 |
| Yes | 1 | | |
| No. of antihypertensive drugs: | | | |
| 1 | 6.98 | | 0.00 |
| 2 | 7.21 | 2.74-17.74 | 0.00 |
| 3-4 | 1 | 2.64-19.72 | 0.00 |
| Smoking: | | | |
| No | 7.46 | 4.19-13.27 | 0.00 |
| Yes | 1 | | |
| Perception of susceptibility: | | | |
| Low | 1 | | |
| High | 2.64 | 1.48-4.69 | 0.00 |
| Perception of barrier: | | | |
| Low | 3.24 | 1.83-5.75 | 0.00 |
| High | 1 | | |
| Cue to action: | | | |
| Low | 1 | | |
| High | 0.34 | 0.18-0.65 | 0.00 |

Table (2): Final regression model of participant's treatment compliance