

	Number	Percent
Negative	5	5%
Positive anticytoplasmic ribosomal antibody	1	1%
Positive homogenous&speckled	1	1%
Positive nucleolar	1	1%
Positive speckled	68	68%
Total	100	100%

Table 1 show ANA pattern

Antibody type	Number	Percent
Nucleosome	20	20%
DsDNA	41	41%
Histones	17	17%
Sm	5	5%
RNP	13	13%
Sm/RNP	14	14%
SS-A/Ro60KD	44	44%
SS-A/Ro52KD	40	40%
SS-B/La	8	8%
Scl70	4	4%
RNA-Polymerase	1	1%
Ku	2	2%
PM-Scl100	3	3%
Mi-2	2	2%
JO-1	3	3%
PCNA	5	5%
Ribosome Po	5	5%
DFS-70	4	4%

Table 2. ANA profile

	B	S.E.	Wald	Sig.	Odd's ratio	95% C.I.for EXP(B)	
						Lower	Upper
age	-.990-	1.352	.536	.464	.372	.026	5.254
ANA	-19.298-	40192.98	.000	0.001	.000	.000	.
Disease duration	20.510	13397.65	.000	.999	8.077E8	.000	.
p/c ratio	17.777	11746.70	.000	.999	52518911.	.000	.
s.creatinine	20.287	40192.97	.000	1.000	6.467E8	.000	.
renalbiopsy	-18.816-	11746.70	.000	.999	.000	.000	.
Hemoptysis	21.896	40192.96	.000	0.001	3.231E9	.000	.
Cough	-21.896-	13397.65	.000	0.001	.000	.000	.
expectoration	-17.561-	17282.72	.000	0.001	.000	.000	.
Dyspnea	3.497	1.015	11.866	0.001	33.000	.000	.
chestpain	.439	23571.33	.000	1.000	1.552	.000	.
fever	21.146	17282.72	.000	0.001	1.527E9	.000	.

**Table 3. Univariate Binary Regression Analysis between
Cases of ILD and controls**