

Table (1): Socio-demographic characteristics and weight of the studied cases (No. =50).

Parameter	Median or Number (Frequency)
Age	
Median (Range)	8 (1-156)
< 1 month	5 (10%)
1 month - < 1 year	25 (50%)
1 -5 years	14 (28%)
> 5 years	6 (12%)
Gender	
Males	34 (68%)
Females	16 (32%)

Table (2): Clinical presentation of the studied cases (No.=50)

Parameter	Number and Frequency
Heart failure	
No	23 (46)
Yes	27 (54)
Dyspnea	
No	22 (44)
Yes	28 (56)
Loss of femoral pulse	
No	18 (36)
Yes	32 (64)
Femoral radial delay	
No	0 (0.0)
Yes	50 (100)
Hypertension	
No	29 (58)
Yes	21 (42)

Table (3): Echocardiographic findings of the studied cases. (No.=50).

Parameter	Summary statistics
Left ventricular hypertrophy	
No	21 (42%)
Yes	29 (58%)
Hypoplasia of aortic arch	
No	44 (88%)
Yes	6 (12%)
Bicuspid aortic valve	
No	39 (78%)
Yes	11 (22%)
Aortic Stenosis	
No	41 (82%)
Yes	9 (18%)
Doppler gradient at the coarctation	
Mean± S.D.	48.5 ± 9.7
Median (Range)	45.5 (35– 70)

Table (4): Distribution of the studied cases according to associated cardiac congenital anomalies (No.=50).

Parameter	Number and Frequency
ASD	7 (14)
VSD	7 (14)
PDA	6 (12)
PFO	1 (2)
TGA	1 (2)
TR	1 (2)
MS	1 (2)
Complete A-V canal defect	1 (2)
PDA, ASD	1 (2)
PDA, VSD	1 (2)
Shone complex	1 (2)
PDA, VSD, DORV	1 (2)