

Table 1: Effect of the tested agents on ALT, AST, AST/ALT, TC and TG levels level against dexamethasone-induced NAFLD in rats

	Control	Dexamethason	Rosuvastatin	Stevia	Rosuvastatin + Stevia
ALT (U/L)	18.17 ± 1.08	56.17 ± 2.54 ^a	21.33 ± 1.54 ^{bc}	28.17 ± 0.91 ^{ab}	20.00 ± 1.57 ^{bc}
AST (U/L)	14.67 ± 1.33	54.50 ± 2.84 ^a	25.00 ± 2.27 ^{ab}	28.67 ± 1.71 ^{ab}	21.17 ± 1.60 ^{ab}
AST/ALT	0.82 ± 0.08	0.98 ± 0.08	1.21 ± 0.14	0.99 ± 0.06	1.08 ± 0.12
TC (mg/dl)	88.50 ± 7.92	197.70 ± 13.17 ^a	110.50 ± 6.47 ^b	115.5 ± 10.81 ^b	84.17 ± 7.86 ^b
TG (mg/dl)	61.95 ± 5.74	163.2 ± 11.72 ^a	96.02 ± 4.90 ^{ab}	101.8 ± 6.20 ^{ab}	65.78 ± 5.21 ^{bcd}

Data represent mean ± SEM of six observations. ^aSignificant variation at p <0.01 vs. Control group.

^bSignificant variation at p <0.0001 vs. DEX group. ^cSignificant variation at p <0.05 vs. Stevia group.

^dSignificant variation at p <0.05 vs. Rosuvastatin group.