

Variables	During attack	After remission	P-value
Serum creatinine Mean \pm SD	0.39 \pm 0.12	0.41 \pm 0.08	0.25
Serum urea Mean \pm SD	20.16 \pm 7.23	18.53 \pm 2.07	0.29
Serum Albumin Mean \pm SD	1.97 \pm 0.53	3.32 \pm 0.34	<0.001
total serum proteins Mean \pm SD	4.42 \pm 0.89	7.02 \pm 0.35	<0.001
Serum cholesterol Mean \pm SD	394.82 \pm 119.67	191.45 \pm 21.75	<0.001
Serum triglycerides Mean \pm SD	244.63 \pm 79.27	134.71 \pm 26.47	<0.001
24hr urinary proteins Mean \pm SD	5673.16 \pm 5402.15	196.51 \pm 48.60	<0.001
Albumin-to-globulin ratio Mean \pm SD	0.76 \pm 0.29	0.71 \pm 0.15	0.31
TSH Mean \pm SD	5.75 \pm 4.52	4.27 \pm 3.31	<0.001
TT3 Mean \pm SD	112.02 \pm 43.92	141.52 \pm 42.82	<0.001
TT4 Mean \pm SD	5.86 \pm 2.20	8.13 \pm 2.21	<0.001

Table (1): The biochemical markers in children with nephrotic syndrome during the attack and after remission.

Thyroid status	During attack	After remission
Euthyroid	15 (29.41%)	36 (70.59%)
Subclinical hypothyroidism	24 (47.06%)	14(27.45%)
True hypothyroidism	12 (23.52%)	1 (1.96%)

Table (2): Thyroid status in the study participants during the attack and after remission.