

		<i>RF Group</i> (n = 29)	<i>IA Group</i> (n = 29)	95%	p
Age (years)		45.31 ± 8.85	44.76 ± 9.88	- 4.38, 5.49	0.82
Gender	Male	35% (10)	45% (13)	-0.15, 0.35	0.59
	Female	65% (19)	55% (16)		
BMI		27 ± 5.86	24.86 ± 7.07	-1.28, 5.56	0.22
ASA	I	86% (25)	93% (27)	-0.09, 0.23	0.67
	II	14% (4)	7% (2)		
	II	38% (11)	45% (13)	-0.18, 0.32	0.79
	III	62% (18)	55% (16)		
Osteoarthritis Side	Right	59% (17)	59% (17)	-	1
	Left	41% (12)	41% (12)		
Duration (years)		4.48 ± 1.18	4.1 ± 0.9	-0.17, 0.93	0.18

Table (1): Demographic characteristics of the patients.

<i>WOMAC score</i>	<i>RF Group (n = 29)</i>	<i>IA Group (n = 29)</i>	<i>95% CI</i>	<i>p</i>
WOMAC PAIN	Basal	11.72 ± 1.907	11.34 ± 2.159	-0.69, 1.45
	Two weeks	7.48 ± 1.617	7.52 ± 1.214	-0.79, 0.72
	One month	5.90 ± 1.012	7.00 ± 1.512	-1.78, -0.43
	Two months	5.83 ± 1.560	7.17 ± 1.713	-2.21, -0.48
	Three months	7.00 ± 1.982	8.07 ± 2.329	-2.21, 0.07
	Six months	7.86 ± 2.295	8.90 ± 2.396	-2.27, 0.20
WOMAC Stiffness	Basal	2.62 ± 0.942	2.55 ± 1.152	-0.48, 0.62
	Two weeks	0.41 ± 0.628	0.69 ± 0.930	-0.69, 0.14
	One month	0.28 ± 0.455	0.45 ± 0.632	-0.46, 0.12
	Two months	0.14 ± 0.351	0.41 ± 0.628	-0.54, -0.01
	Three months	0.55 ± 0.632	0.72 ± 0.797	-0.55, 0.21
	Six months	1.03 ± 0.680	1.07 ± 0.961	-0.47, 0.40
WOMAC Function	Basal	40.76 ± 6.004	40.21 ± 3.144	-1.97, 3.07
	Two weeks	33.97 ± 4.395	34.52 ± 3.612	-2.67, 1.56
	One month	26.90 ± 4.135	27.38 ± 3.590	-2.52, 1.55
	Two months	23.28 ± 3.217	25.31 ± 2.316	-3.51, -0.56
	Three months	28.28 ± 3.239	32.48 ± 3.562	-6.00, -2.42
	Six months	32.59 ± 4.005	35.31 ± 4.630	-5.00, -0.45
WOMAC Total	Basal	54.72 ± 5.812	53.86 ± 5.242	-2.05, 3.77
	Two weeks	42.24 ± 5.356	42.72 ± 4.008	-2.97, 2.01
	One month	33.83 ± 4.310	34.83 ± 4.132	-3.22, 1.22
	Two months	28.14 ± 4.397	33.93 ± 4.464	-8.12, -3.46
	Three months	35.17 ± 4.833	40.48 ± 6.801	-8.41, -2.21
	Six months	41.07 ± 4.114	46.17 ± 5.478	-7.65, -2.56

Table (2): WOMAC score of the patients.