



Comparison of intraarticular injection versus radiofrequency neurotomy in knee osteoarthritis.

Ahmed Talat Mohamed Ahmed Elsaied Abdelrahman, Fawzy
A. Badawy, Wael Alham Mahmoud,.

Department of anesthesia, faculty of medicine, Sohag university

Abstract

Background: Multiple modalities are existing for pain management in knee osteoarthritis cases. This study was conducted to compare the efficacy of genicular nerves radiofrequency ablation versus intraarticular steroid injection in pain management in knee osteoarthritis.

Patients and methods: A total of 58 cases with knee osteoarthritis were included and they were divided into 2 equal groups; the RF group who underwent radiofrequency for the genicular nerves, and the IA group who underwent intraarticular steroid injection. Follow up visits were scheduled after 1 week, 2 weeks, 1, 2, 3, and 6 months. Both visual analog score and The Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) scores were assessed in each visit. Furthermore, patient satisfaction was also recorded.

Results: Patient characteristics did not differ significantly between the two groups ($p > 0.05$). Both VAS and total WOMAC scores were improved in both groups. However, the RF showed better scores at 2-, 3-, and 6- month visits. Satisfaction grades did not differ between the two groups.

Conclusion: Both genicular nerve RF and intraarticular steroid injection are safe and efficacious in pain management in knee osteoarthritis cases. Nevertheless, the effect is more prolonged after RF.

Keywords: intra-articular, osteoarthritis, radiofrequency

Introduction

Knee osteoarthritis is one of the commonest joint diseases that is characterized by progressive joint cartilage degeneration. Pain is the most annoying symptom as reported by such cases (1).

Therefore, the main goal of osteoarthritis medications is to relieve pain. Other goals include an increased range of motion, decreasing disability, and protecting the joint from further damage (2).

Multiple treatment modalities are present for knee osteoarthritis including weight loss, non-steroidal anti-inflammatory drugs, glucosamine, intraart-

icular injections, arthroscopy, and surgical intervention (3).

Intraarticular injection techniques are usually recommended before surgical interventions. Multiple studies have reported that intraarticular steroid injection can improve knee pain, function, and range of motion. Steroid injections have been also beneficial in cases with exacerbation of joint swelling or inflammation as they are potent anti-inflammatory drugs (4).

Radiofrequency neurotomy has been successfully used in the management of many painful conditions like trigeminal neuralgia and spinal pain (5).

Furthermore, one study has reported that radiofrequency genicular nerve ablation may provide adequate and prolonged pain relief in patients of chronic osteoarthritis (6).

Genicular nerves supplying the knee joint arise from obturator, femoral, saphenous, common peroneal, and tibial nerves. Application of RF causes destruction of pain signals leading to interruption of pain signaling pathways (7) This study was conducted aiming to compare the efficacy of genicular nerve radiofrequency ablation and intraarticular steroid injection in knee osteoarthritis.

Patients and methods:

Study design: This is a prospective comparative single-blinded study, that was conducted at Sohag University Hospitals during the period between January and July 2019.

Study cases: A total of 58 cases who presented to the outpatient pain clinic complaining of knee osteoarthritis were enrolled in the study. They were divided into 2 equal groups; the RF group included 29 cases who underwent genicular nerve radiofrequency ablation, and the IA group included the remaining 29 cases who underwent intraarticular steroid injection.

Randomization: was done through a computer program.

Ethical considerations: approval of ethics committee at Sohag University then a written consent from each patient after full illustration of the procedure.

Inclusion criteria: Patients who were diagnosed with knee osteoarthritis according to the American College of Rheumatology (8), and failed to respond to conventional treatment techniques including medications and physiotherapy. Patients with Kallgren-Lawrence (K/L) rating scale (9) lower than 4 were included in our study.

Exclusion criteria: Patients with acute

knee pain, previous knee surgeries, bleeding diathesis, connective tissue disorders, or neurological disabilities were excluded from our study. Furthermore, patients with a Kallgren-Lawrence (K/L) rating scale of 4 were also rolled out.

Patient consent:

Informed written consent was obtained before the procedure from all cases after explaining the advantages and drawbacks of each approach.

Pre-procedure preparation:

All cases were subjected to complete history taking, physical examination, and routine laboratory investigations. Additionally, plain X-ray films were ordered for all cases. Both visual analog score (VAS) and Western Ontario and McMaster Universities Osteoarthritis (WO-MAC) (10) scores were assessed before the procedure in all cases.

Intraarticular injection:

The medial approach was the preferred method of injection when the patient was in the supine position. Methylprednisolone (MP) was administered via intraarticular injection at a dose of 8mg/mL×5mL.

Radiofrequency genicular nerve ablation:

The patient was placed in the supine position and the possible locations of the genicular nerves were identified under fluoroscopic guidance on the medial and lateral aspects of the lower femur in addition to the medial aspect of the upper tibia. After skin sterilization and draping, it was infiltrated with lidocaine (2% concentration), 5 ml volume at first. After that, a 10 cm RF cannula with a 22-gauge size was placed for each nerve. Multiple images in different planes were obtained to confirm the needle tip position. The electrode tip temperature was raised up to 80° C for 1.5 minutes.

Follow-up: Scheduled follow-up visits were arranged after 1, and 2 weeks away from the procedure. Patients were examined for potential complications like infection or hematoma. Additional 4 visits were scheduled at 1,2,3 and 6 months after the intervention. Both VAS and WOMAC scores were assessed during these visits. Moreover, patient satisfaction was also recorded.

Sample size calculation: Sample size was calculated using Power Analysis and Sample Size software program (PASS) version 15.0.5 for windows (2017) using the results published by S. Sari et al (August 2016) with the WOMAC score difference during a three-month follow-up period between both groups after the injection as the primary outcome. Patients were allocated into two groups: Group RF: patients received an intra-articular radiofrequency neurotomy ablation and Group IA: patients received an intra-articular steroid injection. The null hypothesis was considered as the absence of difference between treatment modalities regarding the WOMAC score difference during a three-month follow-up period of the injection. A sample size of 23 patients in each group is needed to achieve 90% power ($1-\beta$ or the probability of

rejecting the null hypothesis when it is false) in the proposed study using a Repeated Measures Analysis with a significance level (α or the probability of rejecting the null hypothesis when it is true) of 5%. The expected drop-outs are 6 so a total of 29 will be enrolled in each group.

Statistical analysis:

Statistical analysis was performed using the SPSS software program. Data were presented as mean \pm standard deviation, or number (%). The comparison between both groups was performed by an independent t-test or chi-squared test as appropriate. Changes were evaluated using the repeated measures general linear model. A p-value of less than 0.05 was considered statistically significant.

Results

The mean age of the included cases was 45.31 and 44.76 years for RF and IA groups respectively. We included 19 females (65%) and 10 males (35%) in the RF group, while the IA group included 16 females (55%) in addition to 13 males (45%). The rest of the patient characteristics are shown in table (1). Neither of these variables was significantly different between the two groups.

		RF Group (n = 29)	IA Group (n = 29)	95%	p
Age (years)		45.31 \pm 8.85	44.76 \pm 9.88	- 4.38, 5.49	0.82
Gender	Male	35% (10)	45% (13)	-0.15, 0.35	0.59
	Female	65% (19)	55% (16)		
BMI		27 \pm 5.86	24.86 \pm 7.07	-1.28, 5.56	0.22
ASA	I	86% (25)	93% (27)	-0.09, 0.23	0.67
	II	14% (4)	7% (2)		
	II	38% (11)	45% (13)	-0.18, 0.32	0.79
	III	62% (18)	55% (16)		
Osteoarthrit is Side	Right	59% (17)	59% (17)	-	1
	Left	41% (12)	41% (12)		
Duration (years)		4.48 \pm 1.18	4.1 \pm 0.9	-0.17, 0.93	0.18

Table (1): Demographic characteristics of the patients.

In both groups, the total WOMAC score was decreased compared to the baseline score. Nevertheless, the RF group showed significantly better scores at 2-,3-, and 6- month visits

compared to the steroid group, indicating that RF may have a prolonged duration of pain relief compared to steroids ($p < 0.001$). These data are illustrated in table (2).

WOMAC score		RF Group (n = 29)	IA Group (n = 29)	95% CI	p
WOMAC PAIN	Basal	11.72 ± 1.907	11.34 ± 2.159	- 0.69, 1.45	0.444
	Two weeks	7.48 ± 1.617	7.52 ± 1.214	-0.79, 0.72	0.927
	One month	5.90 ± 1.012	7.00 ± 1.512	- 1.78, - 0.43	0.003
	Two months	5.83 ± 1.560	7.17 ± 1.713	-2.21, -0.48	0.003
	Three months	7.00 ± 1.982	8.07 ± 2.329	-2.21, 0.07	0.114
	Six months	7.86 ± 2.295	8.90 ± 2.396	-2.27, 0.20	0.080
WOMAC Stiffness	Basal	2.62 ± 0.942	2.55 ± 1.152	-0.48, 0.62	0.840
	Two weeks	0.41 ± 0.628	0.69 ± 0.930	-0.69, 0.14	0.305
	One month	0.28 ± 0.455	0.45 ± 0.632	-0.46, 0.12	0.328
	Two months	0.14 ± 0.351	0.41 ± 0.628	-0.54, -0.01	0.057
	Three months	0.55 ± 0.632	0.72 ± 0.797	-0.55, 0.21	0.470
	Six months	1.03 ± 0.680	1.07 ± 0.961	-0.47, 0.40	0.980
WOMAC Function	Basal	40.76 ± 6.004	40.21 ± 3.144	-1.97, 3.07	0.488
	Two weeks	33.97 ± 4.395	34.52 ± 3.612	-2.67, 1.56	0.601
	One month	26.90 ± 4.135	27.38 ± 3.590	-2.52, 1.55	0.637
	Two months	23.28 ± 3.217	25.31 ± 2.316	-3.51, -0.56	0.002
	Three months	28.28 ± 3.239	32.48 ± 3.562	-6.00, -2.42	< 0.001
	Six months	32.59 ± 4.005	35.31 ± 4.630	-5.00, -0.45	0.008
WOMAC Total	Basal	54.72 ± 5.812	53.86 ± 5.242	-2.05, 3.77	0.350
	Two weeks	42.24 ± 5.356	42.72 ± 4.008	-2.97, 2.01	0.699
	One month	33.83 ± 4.310	34.83 ± 4.132	-3.22, 1.22	0.371
	Two months	28.14 ± 4.397	33.93 ± 4.464	-8.12, -3.46	< 0.001
	Three months	35.17 ± 4.833	40.48 ± 6.801	-8.41, -2.21	< 0.001
	Six months	41.07 ± 4.114	46.17 ± 5.478	-7.65, -2.56	< 0.001

Table (2): WOMAC score of the patients.

Although VAS scores were decreased compared to the baseline in both groups, the RF group showed significantly lower VAS scores in the last 3 follow up visits ($p < 0.05$). Based on VAS and WOMAC score findings, it was expected to see better patient satisfaction

during the last 3 follow up visits. However, patient satisfaction did not differ between the two groups. This may be attributed to having better scores compare to the baseline which may have a positive impact in the patient daily life.




Figure (1):
Total
WOMAC
score changes
through the
study period.




Figure (2): VAS
score changes
through the study
period.

Discussion

Osteoarthritis is the most common chronic joint disease with worldwide prevalence. Its manifestations include pain, joint dysfunction, and muscle atrophy. The knee joint, as it is a weight-bearing joint, is the commonest joint

affected by that disease (11). As the main pathophysiological process of OA cannot be reversed, the existing treatment modalities including lifestyle changes, physical therapy, medications, and surgery are used for symptom con-

trol (12). RF can be applied for knee osteoarthritis in two ways; either extra-articular ablation of genicular nerves, or intraarticular pulsed radiofrequency (6, 13). This study was conducted at Suhag University hospitals aiming to evaluate the safety and efficacy of genicular nerve radiofrequency ablation versus intraarticular steroid injection in knee osteoarthritis patients. Unluckily, there is a paucity of studies comparing RF neurotomy and intraarticular injection. We included a total of 58 cases, and they were divided into 2 equal groups; the RF group who underwent genicular nerve RF, and the IA group who underwent steroid injection. No statistically significant difference was detected between the two groups regarding patient demographics (age, sex, BMI, and OA grade) ($p > 0.05$).

Another study handling the same perspective did not report any statistically significant difference between patient characteristics before intervention (4). This comes in line with our study results. Our study revealed that both groups experienced a decrease in both VAS and WOMAC scores after both procedures. However, the radiofrequency group had better scores compared to steroids especially after 2, 3, and 6 months ($p < 0.05$). Another study has also compared RF with steroid injection in managing knee pain. Authors reported results that are consistent with ours. The RF group reported a significant reduction regarding both VAS and WOMAC scores at the first month and the third month ($p < 0.001$) (4). Conversely, another study reported no difference between the two approaches regarding pain and function improvement at 3 and 6 months after knee arthroplasty (15). In another study, Our study has some drawbacks, first of all, is relatively small sample size, and short duration follow up. In addition, the analgesic requirement of

the patients was not assessed as well. These points must be kept in consideration on performing further studies.

Conclusion:

Both genicular nerve RF and intraarticular steroid injection are safe and efficacious in pain management in knee osteoarthritis cases. Nevertheless, the effect is more prolonged after RF.

References:

1. Sampath S, Harshavardhan JG. Efficacy of intraarticular steroid injection in knee osteoarthritis using knee society score and visual analog score. *International Journal of Research in Orthopaedics*. 2019;5(2):299.
2. del Carmen Caamaño M, García-Padilla S, Duarte-Vázquez MÁ, González-Romero KE, Rosado JL. A double-blind, active-controlled clinical trial of sodium bicarbonate and calcium gluconate in the treatment of bilateral osteoarthritis of the knee. *Clinical Medicine Insights: Arthritis and Musculoskeletal Disorders*. 2017;10:1179544116688899.
3. Bruyère O, Cooper C, Pelletier J-P, Maheu E, Rannou F, Branco J, et al., editors. A consensus statement on the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO) algorithm for the management of knee osteoarthritis—from evidence-based medicine to the real-life setting. *Seminars in arthritis and rheumatism*; 2016: Elsevier.
4. Bruyère O, Cooper C, Cutolo M, Reginster J-Y. International endorsement of the ESCEO algorithm for management of knee osteoarthritis in clinical practice. *Semin Arthritis Rheum*. 2017;47(2):e10.
5. Sarı S, Aydın ON, Turan Y, Özlülerden P, Efe U, Kurt Ömürlü İ. Which one is more effective for the clinical treatment of chronic pain in knee osteoarthritis: radiofrequency neurotomy of the genicular nerves or intra-articular injection? *International journal of rheumatic diseases*. 2018;21(10):1772-8.

6. Masala S, Fiori R, Raguso M, Morini M, Calabria E, Simonetti G. Pulse- dose radiofrequency for knee osteoarthritis. *Cardiovascular and interventional radiology*. 2014;37(2):482-7.
7. Choi W-J, Hwang S-J, Song J-G, Leem J-G, Kang Y-U, Park P-H, et al. Radiofrequency treatment relieves chronic knee osteoarthritis pain: a double- blind randomized controlled trial. *PAIN®*. 2011;152(3):481-7.
8. Le PU. Is genicular nerve radiofrequency ablation safe? A literature review and anatomical study. *Pain Physician*. 2016;19:E697-E705.
9. Altman RD. Criteria for classification of clinical osteoarthritis. *The Journal of rheumatology Supplement*. 1991;27:10-2.
10. Kellgren J, Lawrence J. Radiological assessment of osteoarthrosis. *Annals of the rheumatic diseases*. 1957;16(4):494.
11. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *The Journal of rheumatology*. 1988;15(12):1833-40.
12. Sakellariou G, Conaghan PG, Zhang W, Bijlsma JW, Boyesen P, D'agostino MA, et al. EULAR recommendations for the use of imaging in the clinical management of peripheral joint osteoarthritis. *Annals of the rheumatic diseases*. 2017;76(9):1484-94.
13. Vaishya R, Pariyo GB, Agarwal AK, Vijay V. Non-operative management of osteoarthritis of the knee joint. *Journal of clinical orthopedics and trauma*. 2016;7(3):170-6.
14. Karaman H, Tüfek A, Kavak GÖ, Yildirim ZB, Uysal E, Celik F, et al. Intra-articularly applied pulsed radiofrequency can reduce chronic knee pain in patients with osteoarthritis. *Journal of the Chinese Medical Association*. 2011;74(8):336-40.
15. Takahashi K, Kurosaki H, Hashimoto S, Takenouchi K, Kamada T, Nakamura H. The effects of radiofrequency hyperthermia on pain and function in patients with knee osteoarthritis: a preliminary report. *Journal of Orthopaedic Science*. 2011;16(4):376-81.
16. Qudsi-Sinclair S, Borrás-Rubio E, Abellan-Guillén JF, Padilla del Rey ML, Ruiz-Merino G. A Comparison of Genicular Nerve Treatment Using Either Radiofrequency or Analgesic Block with Corticosteroid for Pain after a Total Knee Arthroplasty: A Double-Blind, Randomized Clinical Study. *Pain Practice*. 2017;17(5):578-88.
17. Ikeuchi M, Ushida T, Izumi M, Tani T. Percutaneous radiofrequency treatment for refractory anteromedial pain of osteoarthritic knees. *Pain Medicine*. 2011;12(4):546-51.
18. Wong J, Bremer N, Weyker PD, Webb CA. Ultrasound-guided genicular nerve thermal radiofrequency ablation for chronic knee pain. *Case reports in anesthesiology*. 2016;2016.
19. Ferdinand Iannaccone D, Samuel Dixon R, Andrew Kaufman M. A review of long-term pain relief after genicular nerve radiofrequency ablation in chronic knee osteoarthritis. *Pain physician*. 2017;20:E437-E44.
20. Mogahed M, Mohamed R, Refaat HM. Intraarticular Pulsed Radiofrequency vs. Radiofrequency Neurotomy in Patients with Chronic Knee Pain due to Osteoarthritis (OA). *J Anesth Clin Res*. 2017;8(10):2-6.
21. Bellini M, Barbieri M. Cooled radiofrequency system relieves chronic knee osteoarthritis pain: the first case-series. *Anesthesiology intensive therapy*. 2015;47(1):30-3.
22. Arroll B, Goodyear-Smith F. Corticosteroid injections for osteoarthritis of the knee: a meta-analysis. *BMJ*. 2004;328(7444):869.

