

Table-1. 2019 EULAR/ACR Classification criteria for systemic lupus erythematosus (SLE) (11).

Entry criterion			
Anti nuclear antibodies (ANA) at a titer of $\geq 1:80$ on HEp-2 cells or an equivalent positive test (ever)			
↓			
If absent, do not classify as SLE If present, apply additive criteria			
↓			
Additive criteria			
Do not count a criterion if there is a more likely explanation than SLE			
Occurrence of a criterion on at least one occasion is sufficient. SLE classification requires at least one clinical criterion and ≥ 10 points			
Criteria need not occur simultaneously			
Within each domain, only the highest weighted criterion is counted toward the total scores			
Clinical domains	Weight	Immunology domains	Weight
Constitutional		Antiphospholipid antibodies	
Fever	2	Anti-cardiolipin antibodies OR Anti-B2GP1 antibodies OR	
Hematologic		lupus anticoagulant	2
Leukopenia	3		
Thrombocytopenia	4		
Autoimmune hemolysis	4		
Neuropsychiatric		Complement proteins	
Delirium	3	Low C3 OR low C4	3
Psychosis	3	Low C3 AND low C4	4
Seizure	5	SLE-specific antibodies	
		Anti-dsDNA antibody OR Anti-Smith antibody	6
Mucocutaneous			
Non-scarring alopecia	2		
Oral ulcers	2		
Subacute cutaneous OR discoid lupus	4		
Acute cutaneous lupus	6		
Acute cutaneous lupus	6		

Serosal		
Pleural or pericardial effusion	5	
Acute pericarditis	6	
Musculoskeletal		
Joint involvement	6	
Renal		
Proteinuria > 0.5 g/24h	4	
Renal biopsy Class II or V lupus nephritis	8	
Renal biopsy Class III or IV lupus nephritis	8	
Total score:		



Classify as SLE with a score of 10 or more if entry criterion fulfilled

Table-2 Classification of lupus nephritis (15).

ISN/RPS class	Description
I	Minimal Mesangial Lupus Nephritis
II	Mesangial proliferative Lupus Nephritis
III	Focal Lupus Nephritis
IV	Diffuse Lupus Nephritis
V	Membranous Lupus Nephritis
VI	Advanced Sclerosing Lupus Nephritis