

Table (1):Distribution of the studied patients according to sensitivity pattern to different antibiotics.

Antibiotic	Sensitive NO. (%)	Resistant NO. (%)
Cefoxitin	11(11.0%)	89(89.0%)
PenicillinG	0(0.0%)	100(100%)
Oxacillin	11(11.0%)	89(89.0%)
Gentamycin(CN)	66(66.0%)	34(34.0%)
Ciprofloxacin	74(74.0%)	26(26.0%)
Levofloxacin	74(74.0%)	26(26.0%)
Moxifloxacin	96(96.0%)	4(4.0%)
Erythromycin	56(56.0%)	44(44.0%)
Clindamycin	62(62.0%)	38(38.0%)
Quinipristin/Dalfopristin	100(100%)	0(0%)
Linezolid	100(100%)	0(0%)
Vancomycin	100(100%)	0(0%)
Tetracycline	42(42.0%)	58(58.0%)
Tigecycline	100(100%)	0(0%)
Nitrofurantoin	100(100%)	0(0%)
Rifampicin	100(100%)	0(0%)
Trimethoprim/Sulfamethoxazole	83(83.0%)	17(17.0%)
Teicoplanin	100(100%)	0(0%)
Doxycycline	42(42.0%)	58(58.0%)
Azithromycin	89(89.0%)	11(11.0%)
Chloramphenicol	56(56.0%)	44(4.0%)