

Table (1): Percentage change in PT measurement over time of studied population in control group.

PT	At room temperature	Number (%) of samples with >10% change	At 4 C°	Number (%) of samples with >10% change
At 4 hours Mean ± SD Median (range)	-1.07±4.03 -0.89 (-8.45-8.00)	0	1.07±6.09 0.89 (-16.26-12.12)	3 (6.98%)
At 8 hours Mean ± SD Median (range)	-1.36±4.29 -0.79 (-11.38-5.74)	2 (4.65%)	1.53±6.65 2.65 (-16.26-11.81)	5 (11.63%)
At 24 hours Mean ± SD Median (range)	1.72±3.78 0.96 (-3.82-10.08)	1 (2.33%)	2.08±10.45 2.33 (-30.08-36.61)	8 (18.60%)
Acceptable time	24 hours		24 hours	

Table (2): Percentage change in PT measurement over time of studied population in Anticoagulant group.

PT	At room temperature	Number (%) of samples with >10% change	At 4 C°	Number (%) of samples with >10% change
At 4 hours Mean ± SD Median (range)	2.60±5.81 0.73 (-9.47:20.14)	4 (9.30%)	0.84±4.03 0.66 (-8.38:12.35)	2 (4.65%)
At 8 hours Mean ± SD Median (range)	3.55±4.21 3.31 (-10.53:12.28)	3 (6.98%)	5.29±4.45 4.65 (-8.42:15.07)	4 (9.30%)
At 24 hours Mean ± SD Median (range)	27.39±51.40 14 (1.24:303.68)	33 (76.74%)	19.10±9.80 17.39 (1.86:52.72)	36 (83.72%)
Acceptable time	8 hours		8 hours	