

Table (1): Comparison between group I & II as regard language parameters:

		<i>Mean ± SD</i>		<i>p</i>
		<i>Group I</i>	<i>Group II</i>	
Receptive semantics	Pre-therapy	0.3 ± 0.3	1.05 ± 0.55	0.25
	3m	4.2 ± 1.7	8.95 ± 1.72	0.06
	6m	10.8 ± 3.99	+19.8 ± 3.1	0.09
	9m	19.3 ± 4.6	26.7 ± 3.37	0.21
Expressive semantics	Pre-therapy	0.3 ± 0.3	0.65 ± 0.51	0.56
	3m	1.6 ± 1.28	4.55 ± 1.8	0.19
	6m	4 ± 2.65	8.8 ± 2.7	0.22

		<i>Mean ± SD</i>		<i>p</i>
		<i>Group I</i>	<i>Group I</i>	
Vocabulary size	Pre-therapy	0.7 ± 0.52	1.95 ± 0.81	0.20
	3m	4.5 ± 2.92	11.4 ± 3.79	0.16
	6m	14.9 ± 9.6	22.85 ± 6.69	0.51
	9m	39.4 ± 23.79	48.4 ± 13.91	0.75
Word class	Pre-therapy	1 ± 0	1 ± 0	>0.99
	3m	1 ± 0	1.41 ± 0.19	0.54
	6m	1.5 ± 0.34	1.42 ± 0.16	>0.99
	9m	1.5 ± 0.31	2 ± 0.35	0.35

* = Significant, ** = Highly significant