

Table 1 Preoperative data of studied population

Variables	Group (A): without ILM peeling	Group (B): with ILM peeling	P value
Age/years			
Mean \pm SD	50.07 \pm 10.87	56.67 \pm 9.09	0.08
Median (range)	50 (25-70)	60 (40-67)	
Gender			0.14
Females	5 (33.33%)	9 (60.00%)	
Males	10 (66.67%)	6 (40.00%)	

Table 2: Comparison between the two groups as regard extend of RD

Variables	Group (A): without ILM peeling	Group (B): with ILM peeling	P value
Extend of RD			0.30
One quadrant	1 (6.67%)	0	
Two quadrants	7 (46.67%)	6 (40.00%)	
Three quadrants	3 (20.00%)	4 (26.67%)	
Four quadrants	4 (26.67%)	5 (33.33%)	

Table 3: Comparison between the two groups as regard logMAR BCVA

Variables	Group (A): without ILM peeling	Group (B): with ILM peeling	P value
visual acuity after SOR			0.12
Mean \pm SD	1.18 \pm 0.29	0.99 \pm 0.38	
Median (range)	1.3 (0.6-1.5)	1.1 (0.3-1.5)	0.76
0.3	0	2 (13.33%)	
0.6	1 (6.67%)	1 (6.67%)	
0.8	1 (6.67%)	2 (13.33%)	
0.9	2 (13.33%)	1 (6.67%)	
1	1 (6.67%)	1 (6.67%)	
1.1	1 (6.67%)	3 (20.00%)	
1.2	1 (6.67%)	1 (6.67%)	
1.3	4 (26.67%)	2 (13.33%)	
1.5	4 (26.67%)	2 (13.33%)	