

Table(1): NSAIDs (Ibuprofen) Drug consumption in study groups.

	Group 1 (steroid group)	Group 2(PRF group)	P-value
Basal	600 (200)	600 (200)	
1week	550 (200)	400 (150)	0.01*
1month	550 (200)	400 (200)	0.01*
3month	600 (200)	400 (200)	0.003*

Data presented as mean(SD). * P-value < 0.05

Drug consumption dose expressed in mg per day.




Figure (2): NSAIDs (Ibuprofen) consumption in study groups.

1- Tramadol Drug consumption

There was statistically significant difference in the tramadol consumption post procedure between the two groups specially after one week and one month, steroid group had a decrease in tramadol consumption more then the PRF group (P-value<0.05)

Table(2): Tramadol Drug consumption in study groups.

	Group 1	Group 2	P-value
Basal	250 (100)	250 (100)	
1week	175 (50)	225 (100)	0.005*
1month	175 (50)	225 (100)	0.005*
3month	200 (50)	225 (100)	0.04*

Data presented as mean(SD). * P-value < 0.05

Drug consumption dose expressed in mg per day.




Figure (3): Tramadol consumption in study groups.

Pregabalin Drug consumption

There was no statistically significant difference in the pregabalin consumption post procedure between the two groups, but steroid group had a decrease in pregabalin consumption more than the PRF group (P-value < 0.05)

Table(3): Pregabalin Drug consumption in study groups.

	Group 1	Group 2	P-value
Basal	250 (125)	250 (125)	
1week	175 (75)	225 (100)	0.08
1month	175 (75)	225 (100)	0.08
3month	200 (75)	250 (100)	0.38

Data presented as mean(SD). * P-value < 0.05




Figure (4): pregabalin consumption in study groups.