

Table 1: Clinical and laboratory patients' characteristics (n=100).

Parameter	Mean± SD/ Median (25th-75th) or percentage	Range
History:		
Age (Years)	39 (31-45)	22-55
Smoking (%)	52%	
• Smoking history (years)	14.5 (11.25-20)	5-33
• Cigarette/day	20 (15-30)	10-40
• Brinkman smoking index	300 (202.5-510)	90-1320
Occupation:		
• Manual worker/farmer	53%	
	33%	
	14%	
• Employee		
• Driver		
Sexual history:		
Marriage duration (years)	10 (5-16)	1-26
Patients sexual complaint:		
• Desire problem complaint	7%	
• Erectile dysfunction complaint	57%	
• Premature ejaculation complaint	76%	
• Orgasmic disorder complaint	17%	
• Mixed sexual dysfunction complaint	43%	
Sexual complaint duration (months)	11 (8-18)	6-108
Frequency of intercourse/week	2 (2-3)	1-4
Wife age (years)	33.29±7.47	18-48
Drug abuse history :		
Positive drug abuse history	43%	
Drug abuse duration (years)	5 (3-6.25)	1-20
Route of drug administration:		
• Oral	54.8%	
• Inhalation	23.8%	
• Oral/Inhalation	21.4%	
Types of drug abuse:		
• Tramadol	25%	
• THC	19%	
• Opiates	9%	
• Benzodiazepines	4%	
History of hospital admission due to drug abuse	7.3%	
General examination:		
Weight (Kg)	79.07±10	63-100
Height (M)	1.66± 0.05	1.58-1.80
BMI (Kg/m2)	23.63± 2.37	19.94-27.78
Hormonal profile:		
• Total testosterone (TT) (ng/ml)	6.3 (2.67-7.55)	0.22-9.67
• LH (mIU/ml)	3.1(2.1-5)	0.21-7.13
• FSH (mIU/ml)	3.5(1.86-5.01)	0.23-17.30
Sexual function evaluation:		

SHIM score	15 (12-19)	5-24
EHS	3 (2-3)	1-3
AIPE	19 (16-24)	12-30

Table 2: The result of the binary logistic regression to identify positive drug abuse urine testing using the smoking, history of drug abuse and occupation (driver) variables.

Variable	B	S.E.	Wald	df	Sig.	Exp (B)	95% C.I. for EXP(B)	
							Lower	Upper
Smoking	3.109	.853	13.279	1	.000	22.39	4.207	119.190
History of drug abuse	3.009	.693	18.867	1	.000	20.26	5.213	78.761
Occupation (Driver)	1.018	1.198	.721	1	.396	2.767	.264	28.971

Table (3). Patients classification according the result of drug abuse (n=100).

Parameter	Negative drug abuse n= 65 Mean± SD/ Median (25 th -75 th) or percentage		Positive drug abuse n= 35 Mean± SD/ Median (25 th -75 th) or percentage		P Value
History:					
Age (Years)	40 (31-47)		39 (33-34)		0.553#
Smoking (%)	19%		33%		* <0.0001¶
• Smoking history (years)	16 (10-20)		14 (12-20)		0.894 #
• Cigarette/day	25 (15-40)		20 (15-30)		0.367 #
• Brinkman smoking index	330 (200-560)		300 (205-410)		0.562 #
Occupation:	34%		19%		
• Manual worker/farmer	26%		7%		
• Employee	5%		9%		* 0.018 ¶
• Driver					
Sexual history:					
Marriage duration (years)	11 (4-16)		10 (6-15)		0.983 #
Patients sexual complaint:					
• Desire problem complaint	4%		3%		0.651 ¶
• Erectile dysfunction complaint	36%		21%		0.657 ¶
• Premature ejaculation complaint	50%		26%		0.768 ¶
• Orgasmic disorder complaint	11%		6%		0.978 ¶
• Mixed sexual dysfunction complaint	27%		16%		0.687 ¶
Sexual complaint duration (months)	10 (8-17)		12 (7-20)		0.373 #
Frequency of intercourse/week	2 (2-3)		2 (2-3)		0.825 #
Wife age (years)	33.71± 8		32.51± 6.4		0.419 §
Drug abuse history :					
Positive drug abuse history	12%		31%		* <0.0001¶

Drug abuse duration (yrs)	3 (2.25-4.75)	5 (3-7.75)	0.056 #
Route of drug administration:	9% 3% 0%	14% 7% 9%	0.088 ¶
General examination:			
Weight (Kg)	80.56± 7.65	76.4± 13.93	0.479 §
Height (M)	1.68± 0.03	1.65± 0.08	0.366 §
BMI (Kg/m ²)	23.96± 2.12	23.03± 2.92	0.505 §
Hormonal profile:			
• Total testosterone (TT) (ng/ml)	6.32 (3.08-7.66)	6.24 (2.39-7.4)	0.501 #
• LH (mIU/ml)	3.23 (2.05-5.02)	3.1 (2.1-4.2)	0.963 #
• FSH (mIU/ml)	2.9 (1.87-4.62)	4.21 (1.85-5.6)	0.404 #
Sexual function evaluation:			
SHIM score	15 (11-20)	15 (12-19)	0.742 #
EHS	3 (2-3)	3 (2-3)	0.756 #
AIPE	19 (16-24)	20 (16-25)	0.653 #