

Class	Patients	Percentage
<b>subjective outcome</b>		
Normal A	19	63.33%
Near normal B	8	26.67%
Abnormal C	3	10%
<b>symptoms evaluation</b>		
Normal A	22	73.33%
Near normal B	5	16.67%
Abnormal C	3	10%
<b>Objective outcome after follow up</b>		
Normal A	25	83.33%
Near normal B	3	10%
Abnormal C	2	6.67%
<b>Range of motion of the knee</b>		
Normal A	25	83.33%
Near normal B	3	10%
Abnormal C	2	6.67%
<b>Knee stability</b>		
Normal A	16	53.34%
Near normal B	8	26.66%
Abnormal C	6	20%

**Table 1: Clinical results according to IKDC of the first group.**

Class	Patients	Percentage
<b>subjective outcome</b>		
Normal A	19	63.33%
Near normal B	8	26.67%
Abnormal C	3	10%
<b>symptoms evaluation</b>		
Normal A	22	73.33%
Near normal B	5	16.67%
Abnormal C	3	10%
<b>Objective outcome after follow up</b>		
Normal A	25	83.33%
Near normal B	3	10%
Abnormal C	2	6.67%
<b>Range of motion of the knee</b>		
Normal A	25	83.33%
Near normal B	3	10%
Abnormal C	2	6.67%

<b>Knee stability</b>		
<b>Normal A</b>	<b>16</b>	<b>53.34%</b>
<b>Near normal B</b>	<b>8</b>	<b>26.66%</b>
<b>Abnormal C</b>	<b>6</b>	<b>20%</b>

**Table2: *Clinical results according to IKDC of the second group.***