value (.02).

	20-40 yrs	41-60 yrs	61-80 yrs	>80 yrs
■ Age distribution	2%	31.40%	54.90%	11.70%

Figure (1) and table (1): Age distribution

.

	Farmer	Builder	carpenter	Teacher	House wife	Seller	Police man	Moazen	Plumber	Accountant	Lawyer
■ Occupation	49%	11.80%	9.80%	8.80%	4.90%	4.90%	3.90%	2.90%	2%	1%	1%

Table (3) and figure (3): Patient occupation

Smoking

	Yes	Non
■ Smoking	95.10%	4.90%

Figure (4) and table (4): Smoking incidence

.

Stage grouping

|	Stage I	Stage II	Stage III	Stage IV
■ Stage grouping	13.70%	8.90%	55.90%	21.50%

Figure (5) and Table (5): TNM staging of cancer larynx according to AJCC and UJCC

Degree of differentiation of SCC

|	Well differenciated SCC(G1)	Moderate differenciated SCC (G2)	Poor differenciated SCC(G3)	undifferencia ted SCC (G4)
■ Degree of differentiation of SCC	2%	67.70%	23.50%	3.90%

**Figure (6) and Table (6):** The degree of differentiation of SCC of patients with cancer larynx

Figure (7) Frequency of distant metastasis (M)