

| Pathology | Number of cases | Percentage | Male | Female |
|--|-----------------|-------------|-----------|-----------|
| Hyperplastic lymphoid tissue (Adenoid and other inflammatory results) | 59 | 59% | 30 | 29 |
| Thornwaldt's cyst | 1 | 1% | 0 | 1 |
| Branchial cleft cyst | 1 | 1% | 1 | 0 |
| Lymphoma | 15 | 15% | 7 | 8 |
| Undifferentiated carcinoma | 14 | 14% | 10 | 4 |
| Squamous cell carcinoma | 10 | 10% | 6 | 4 |
| Total | 100 | 100% | 54 | 46 |

Table 1: Illustrates the numbers and percentages of different pathological nasopharyngeal lesions in the whole study population and numbers in each sex group.

| Presenting symptom | Number (percentage %) | Number (percentage %) of benign cases | Number (percentage %) of malignant cases |
|--------------------------|-----------------------|---------------------------------------|--|
| Nasal obstruction | 34 (34%) | 24 (24%) | 10 (10%) |
| Aural symptom | 29 (29%) | 22 (22%) | 7 (7%) |
| Snoring | 19 (19%) | 12 (12%) | 7 (7%) |
| Neck lump | 11 (11%) | 0 (0%) | 11 (11%) |
| Rec epistaxis | 5 (5%) | 2 (2%) | 3 (3%) |
| Dysphagia | 1 (1%) | 0 (0%) | 1 (1%) |
| Headache | 1 (1%) | 1 (1%) | 0 (0%) |
| Total | 100 (100%) | 61 (61%) | 39 (39%) |

Table 2: Illustrates the distribution of patients main complaints in the whole study population and separately in malignant and benign lesions.