

SPSS version 22.0 and Medcalc version 15.8.0, were used for analysis.

<b>Table 1: Patient characteristics:</b>			
<b>Total number</b>	<b>60</b>		
<b>Mean Age by years (SD)</b>	<b>54.83 (10.88)</b>		
<b>Parameter</b>	<b>Yes</b>	<b>No</b>	<b>P-value</b>
<b>Males (%)</b>	<b>43 (71.7%)</b>	<b>17 (28.3%)</b>	<b>0.002</b>
<b>H/O of MI (%)</b>	<b>36 (60%)</b>	<b>24 (40%)</b>	<b>0.001</b>
<b>Dyslipidemia (%)</b>	<b>26 (43.3%)</b>	<b>34 (56.7%)</b>	<b>0.138</b>
<b>Current smoking (%)</b>	<b>25 (41.7%)</b>	<b>35 (58.3%)</b>	<b>0.001</b>
<b>SWMA (%)</b>	<b>33 (55%)</b>	<b>27 (45%)</b>	<b>0.003</b>
<b>Diastolic dysfunction (%)</b>	<b>27 (45%)</b>	<b>33 (55%)</b>	<b>0.001</b>
<b>Mean CIMT &gt; 0.9 mm (%)</b>	<b>24 (40%)</b>	<b>36 (60%)</b>	<b>0.008</b>
<b>ABI &lt; 0.9 mm (%)</b>	<b>20 (33.3%)</b>	<b>40 (66.7%)</b>	<b>0.139</b>
<b>Significant coronary stenosis (&gt;50%) (%)</b>	<b>34 (56.7%)</b>	<b>26 (43.3%)</b>	<b>0.038</b>