

Table (1): Distribution of the studied patients according to sociodemographic criteria (No.=100):

Characteristics	Summary statistics
Gender	
Female	72 (72%)
Male	28 (28 %)
Age at presentation	
Mean± S.D.	27.92 ± 10.52
Median (Range)	25.5 (7 – 60)
Marital status	
Divorced	1 (1%)
Married	54 (54 %)
Single	43 (43%)
Widow	2 (2 %)
Special habits	
No	91 (91%)
Smoker	9 (9%)
Family history	
No	53 (53%)
Yes	47 (47%)

Table (2): Distribution of the studied patients according to disease-related criteria (No.=100).

Characteristics	Summary statistics
Onset	
Acute	33 (33%)
Gradual	59 (59 %)
Insidious	8 (8%)
Course	
Progressive	86 (86%)
Stationary	14 (14%)
General examination	80 (80%)
Free	4 (4%)
Nail affection	12 (12%)
Pallor	2 (2%)
Pallor, nail affection	2 (2%)
Thyroid disease	
The pattern of hair loss	45 (45%)
Androgenetic	14 (14%)
Diffuse	41 (41%)
Localized	
Site of hair loss	8 (8%)
Back	12 (12%)
Diffuse	34 (34%)
Front	21 (21%)
Front, vertex	5 (5%)
Temple	20 (20%)
Vertex	
Scalp examination	2 (2%)
Erythema	88 (88%)
Free	10 (10%)
Scales	
Pull test	74 (74 %)
Negative	26 (26 %)
Positive	
Duration of disease (years)	3.32 ± 3.64
Mean± S.D.	2 (0.5 – 17)
Median (Range)	

Table (3): Distribution of the studied patients according to diagnosis (No.=100).

Diagnosis	Percent
Androgenetic alopecia	45 (45%)
Alopecia areata	26 (26%)
Telogen effluvium	14 (14%)
Tractional alopecia	7 (7%)
Lichen planopilaris	5 (5%)
Tricotillomania	3 (3%)

